

Signatures of doubly charged Higgs in $SU(3)_L \otimes U(1)_N$ model

J. E. Cieza Montalvo¹, Nelson V. Cortez Jr.², M. D. Tonasse³

¹*Instituto de Física, Universidade do Estado do Rio de Janeiro,
Rua São Francisco Xavier 524, 20559-900 Rio de Janeiro, RJ, Brazil*

²*Justino Boschetti 40, 02205-050 São Paulo, SP, Brazil and*

³*Unidade de Registro, Campus Experimental de Registro, Universidade Estadual Paulista,
Rua Tamekishi Takano 5, 11900-000, Registro, SP, Brazil*

(Dated: February 2, 2008)

The scalar sector of the simplest version of the 3-3-1 electroweak models is constructed with three Higgs triplets only. We show that a relation involving two of the constants of the Higgs potential, two vacuum expectation values of the neutral scalars and the mass of the doubly charged Higgs boson leads to important information concerning the signals of this scalar particle.

PACS numbers:

11.15.Ex: Spontaneous breaking of gauge symmetries,

12.60.Fr: Extensions of electroweak Higgs sector,

14.80.Cp: Non-standard-model Higgs bosons.

Doubly charged Higgs bosons are a kind of scalar bosons present in electroweak models which predict scalar triplets with non-zero hipercharge [1]. In the left-right symmetric models, for instance, they are fundamental ingredients for implementation of the seesaw mechanism to generation of neutrino masses [2].

Another important class of electroweak models which predict doubly charged Higgs bosons is the called 3-3-1 models [3, 4]. Here we work with the model version of

the Ref. [4]. In the scalar sector we have only the three scalar triplets $\eta = (\eta^0 \ \eta_1^- \ \eta_2^+)^T$, $\rho = (\rho^- \ \rho^0 \ \rho^{++})^T$ and $\chi = (\chi^- \ \chi^{--} \ \chi^0)^T$, transforming as $(\mathbf{3}, 0)$, $(\mathbf{3}, 1)$ and $(\mathbf{3}, -1)$, respectively, and which are necessary and sufficient to breakdown the symmetry and generate the correct mass spectrum. The most general invariant and renormalizable Higgs potential, obeying $B - L$ discrete symmetry [5], is

$$\begin{aligned} V(\eta, \rho, \chi) = & \mu_1^2 \eta^\dagger \eta + \mu_2^2 \rho^\dagger \rho + \mu_3^2 \chi^\dagger \chi + \lambda_1 (\eta^\dagger \eta)^2 + \lambda_2 (\rho^\dagger \rho)^2 + \lambda_3 (\chi^\dagger \chi)^2 + \\ & + (\eta^\dagger \eta) [\lambda_4 (\rho^\dagger \rho) + \lambda_5 (\chi^\dagger \chi)] + \lambda_6 (\rho^\dagger \rho) (\chi^\dagger \chi) + \lambda_7 (\rho^\dagger \eta) (\eta^\dagger \rho) + \\ & + \lambda_8 (\chi^\dagger \eta) (\eta^\dagger \chi) + \lambda_9 (\rho^\dagger \chi) (\chi^\dagger \rho) + \frac{1}{2} (f \epsilon^{ijk} \eta_i \rho_j \chi_k + \text{H. c.}) \end{aligned} \quad (1)$$

In the potential (1) f and μ_j ($j = 1, 2, 3$) are constants with dimension of mass and the λ_i ($i = 1, \dots, 9$) are adimensional constants [6].

The neutral components of the scalars triplets η , ρ and χ develop non zero vacuum expectation values $\langle \eta^0 \rangle = v_\eta$, $\langle \rho^0 \rangle = v_\rho$ and $\langle \chi^0 \rangle = v_\chi$, with $v_\eta^2 + v_\rho^2 = v_W^2 = (246 \text{ GeV})^2$. The pattern of symmetry breaking is $SU(3)_L \otimes U(1)_N \xrightarrow{\langle \chi \rangle} SU(2)_L \otimes U(1)_Y \xrightarrow{\langle \eta, \rho \rangle} U(1)_{\text{em}}$. From the Higgs potential (1) we can find the masses of the of the neutral scalars

$$m_{H_1^0}^2 \approx 4 \frac{\lambda_2 v_\rho^4 - 2\lambda_1 v_\eta^4}{v_\eta^2 - v_\rho^2}, \quad m_{H_2^0}^2 \approx \frac{v_W^2 v_\chi^2}{2v_\eta v_\rho}, \quad (2a)$$

$$m_{H_3^0}^2 \approx -4\lambda_3 v_\chi^2, \quad m_h^2 = -\frac{f v_\chi}{v_\eta v_\rho} \left[v_W^2 + \left(\frac{v_\eta v_\rho}{v_\chi} \right)^2 \right] \quad (2b)$$

It should be notice that the Eqs. (2b) impose $\lambda_3 < 0$ and $f < 0$. The aproximations in Eqs. (2a) and (2b) are valid for $v_\eta, v_\rho \ll v_\chi$. For the singly charged scalars we

have

$$m_{H_1^\pm}^2 = \frac{v_W^2}{2v_\eta v_\rho} (f v_\chi - 2\lambda_7 v_\eta v_\rho), \quad (2c)$$

$$m_{H_2^\pm}^2 = \frac{v_\eta^2 + v_\chi^2}{2v_\eta v_\chi} (f v_\rho - 2\lambda_8 v_\eta v_\chi) \quad (2d)$$

and for the doubly charged

$$m_{H^{\pm\pm}}^2 = \frac{v_\rho^2 + v_\chi^2}{2v_\rho v_\chi} (f v_\eta - 2\lambda_9 v_\rho v_\chi). \quad (2e)$$

The gauge sector of the 3-3-1 model of the Ref. [4] acommodates one extra Z' neutral boson, two singly charged ($V^\pm$) and two doubly charged ($U^{\pm\pm}$) bosons beyond the

standards Z and $W^\pm$ with mass spectrum

$$m_{Z'}^2 = \frac{2}{3} \left(\frac{e}{s_W} \right)^2 (1 + 3t^2) v_\chi^2, \quad (3)$$

$$m_V^2 = \left(\frac{e}{s_W} \right)^2 \frac{v_\eta^2 + v_\chi^2}{2}, \quad (4)$$

$$m_U^2 = \left(\frac{e}{s_W} \right)^2 \frac{v_\rho^2 + v_\chi^2}{2}, \quad (5)$$

where $s_W = \sin \theta_W$. We have also one heavy lepton (E^+, M^+, T^+) and one heavy quark (J_1, J_2, J_3) in each generation, respectively. The heavy fermions have masses of the order of v_χ . For details on this version of the 3-3-1 models see Ref. [4].

In this work we wish to report about the signals of the doubly charged Higgs when we consider particular values for the adimensional parameter λ_9 in the Higgs potential (1) and for the vacuum expectation value v_χ [6]. The production and signals for the doubly charged Higgs bosons of the 3-3-1 models at LHC collider has been investigated in Ref. [7], where has been found no good signatures for them. However, in this present work we have shown that these signatures are very significant for particular parameter values and due that we are working with masses of order of 500 GeV to 700 GeV, which give better and satisfactory results than that of the previous work. Although we have a reasonable number of free parameters [see Eqs. (1) to (2b)], the model establishes strong bounds among them, these constraints and bounds are really important, bringing good results as can be seen in this manuscript. Throughout this work we take the values of standard model parameters, $m_Z = 91.1876$ GeV, $s_W^2 = 0.23122$, and $m_W = 80.403$ GeV as seen in Ref. [8]. For the 3-3-1 model we take two parameters sets listed in the Table I.

TABLE I: The two parameters sets used in this work. Values of the other variables are given throughout the text.

Set	λ_9	v_η (GeV)	λ_1	λ_2	λ_6	λ_7	λ_8
1	-0.8	195	-1.2	-1	1	-2	-1
2	-1.2						

We fix the general bounds of the adimensional constants of the Higgs potential (1) as $-3 \leq \lambda_i \leq 3$ ($i = 1, \dots, 9$) to guarantee approximately the perturbative regime. Thus, from Eq. (2e) we can see that the inequality $-3 \leq \lambda_9 \leq 0$ is satisfied from the positivity of $m_{H^{\pm\pm}}^2$ because $f \leq 0$.

On the other hand, if we consider $-f \approx v_\chi \gg v_\eta, v_\rho$, we have

$$\lambda_4 \approx 2 \frac{\lambda_2 v_\rho^2 - \lambda_1 v_\eta^2}{v_\eta^2 - v_\rho^2}, \quad \lambda_5 v_\eta^2 + 2\lambda_6 v_\rho^2 \approx -\frac{v_\eta v_\rho}{2} \quad (6)$$

as seen in Ref. [6].




FIG. 1: Parameter f as a function of mass of doubly charged Higgs boson for $\lambda_9 = -0.8$.




FIG. 2: Parameter f as a function of mass of doubly charged Higgs boson for $\lambda_9 = -1.2$.

A simple analysis of Eq. (2e) also shows that the lowest values for λ_9 and f , such to produce values for $m_{H^{\pm\pm}}$ between 100 GeV and 200 GeV, for $v_\chi = 1500$ GeV and $v_\rho = 195$ GeV are $\lambda_9 = -4.50 \times 10^{-3}$, $f = -0.23$ GeV and $\lambda_9 = -1.80 \times 10^{-2}$, $f = -0.92$ GeV, that is, by decreasing the value of λ_9 we increase the value of $m_{H^{\pm\pm}}$.




FIG. 3: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$ for the fermionic sector.




FIG. 4: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$ for the bosonic sector.




FIG. 5: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -1.2$ for the leptonic sector.

with negative f . On the other hand, the values for the parameters λ_5 and λ_6 lead to the constraint $v_\eta > 40.5$ GeV, which can be seen as following from the Eq. (6). In the approximation $-f \approx v_\chi$ it is appropriate to choose the parameter $-1.2 \leq \lambda_9 \leq -0.8$ (see Table II) for the mass of $m_{H^{\pm\pm}} = 500$ GeV. Fig. 1 shows the possible values of f , varying the values of $m_{H^{\pm\pm}}$ for $v_\chi = (1000, 1500, 1700)$ GeV and considering $\lambda_9 = -0.8$ and $v_\eta = 195$ GeV. We can conclude from Fig. 1 that for the $v_\chi = 1000$ GeV we have the acceptable masses up to $m_{H^{\pm\pm}} \simeq 903$ GeV, for $v_\chi = 1500$ GeV we will have up to $m_{H^{\pm\pm}} \simeq 1346$ GeV and for $v_\chi = 1700$ GeV we have consequently up to $m_{H^{\pm\pm}} \simeq 1525$ GeV. If we now consider $\lambda_9 = -1.2$ and the $v_\chi = 1000$ GeV, from Fig. 2 we see that the acceptable masses will be up to $m_{H^{\pm\pm}} \simeq 1107$ GeV, for $v_\chi = 1500$ GeV we will have up to $m_{H^{\pm\pm}} \simeq 1651$ GeV and for $v_\chi = 1700$ GeV we have consequently up to $m_{H^{\pm\pm}} \simeq 1869$ GeV. It should be noticed that the VEV v_η and v_ρ influence very little on the mass of $m_{H^{\pm\pm}}$, so for example for $\lambda_9 = -0.5$, $v_\chi = 1500$ GeV and $v_\eta = 50$ GeV the acceptable mass will be $m_{H^{\pm\pm}} = 1074$ GeV, for $v_\eta = 150$ GeV will be $m_{H^{\pm\pm}} = 1069.5$ GeV and for $v_\eta = 240$ GeV will be $m_{H^{\pm\pm}} = 1061$ GeV. If we take $\lambda_9 = -1.2$, $v_\chi = 1500$ GeV and $v_\eta = 50$ GeV then the acceptable mass will be $m_{H^{\pm\pm}} = 1664$ GeV, for $v_\eta = 150$ GeV we will have $m_{H^{\pm\pm}} = 1656$ GeV and consequently for $v_\eta = 240$ GeV we have $m_{H^{\pm\pm}} = 1644$ GeV, all this happen because in the Eq. (2e) the v_η and v_ρ are related by the relation $v_\eta^2 + v_\rho^2 = (246 \text{ GeV})^2$ and v_ρ is smaller compared with v_χ .

TABLE II: The parameters sets used in this work. Values of the other variables are given throughout the text.

$-f \times 10^3 \text{ GeV}$	0.25	0.88	1.50	2.75	3.99	0.91	1.62	2.55	4.42	6.52
$v_\chi \times 10^3 \text{ GeV}$	1.01	1.01	1.01	1.01	1.01	1.51	1.51	1.51	1.51	1.51
$-\lambda_9$	0.40	0.80	1.20	2.00	2.80	0.50	0.80	1.20	2.00	2.90

Figs. 3 and 4, show the branching ratios for the Higgs decays, $H^{\pm\pm} \rightarrow \text{all}$, where we have chosen for the parameters, masses and the VEV, the following representative values: $\lambda_1 = -1.2$, $\lambda_2 = \lambda_3 = -\lambda_6 = \lambda_8 = -1$, $\lambda_4 = 2.98$, $\lambda_5 = -1.57$, $\lambda_7 = -2$ and $\lambda_9 = -0.8$, $v_\eta = 195$ GeV and $v_\chi = 1500$ GeV. From Fig. 1 we see that the others particles, such as $J_{1,2,3}, T, U^{\pm\pm}, Z', H_2^\pm, H_{2,3}^0$ and h^0 , do not take part in this process because of their masses are larger than the mass of the doubly charged Higgs.

In Figs. 5 and 6 we exhibit the branching ratios for the same particle and for the same parameters, masses and the VEV, as considered in Fig. 3, except that we consider $\lambda_9 = -1.2$. Considering this value, and from

Fig. 2, we see that the acceptable mass for $m_{H^{\pm\pm}}$ is larger and consequently, more particles will take part in this process, as clearly seen from Figs. 5 and 6. Fig. 7 shows the branching ratios: $BR(H^{\pm\pm} \rightarrow V^\pm H_1^\pm)$ and $BR(H^{\pm\pm} \rightarrow H_1^\pm H_2^\pm)$ which vary in a short interval, Fig. 8 shows the cross section for the process $pp \rightarrow H^{++} H^{--}$, which is given in Ref. [7], for the same values of λ 's considered above and $v_\eta = 195$ GeV, except which we put $\lambda_9 = -0.8$ and with other particles masses as given in the Table II, it is to notice that the masses of m_{h^0} , $m_{H_1^\pm}$ and $m_{H_2^\pm}$ depend on the parameter f and therefore they can not be fixed by any value of v_χ .

TABLE III: Values of the masses for $v_\eta = 195$ GeV and the sets of parameters given in the text. All the values in this table are given in GeV. For all values of the parameters for this table we have $m_{H_1^0} = 874$ GeV and $m_{H^\pm} = 348$ GeV.

v_χ	m_E	m_M	m_T	$m_{H_2^0}$	$m_{H_3^0}$	m_V	m_U	$m_{Z'}$	m_{J_1}	m_{J_2}	m_{J_3}
1000	148.9	875	2000	1017.2	2000	467.5	464	1707.6	1000	1410	1410
1500	223.3	1312.5	3000	1525.8	3000	694.1	691.8	2561.3	1500	2115	2115
1700	251.3	1487.5	3400	1729.2	3400	785.2	783.1	2902.8	1700	2397	2397

We have chosen $v_\chi = 1700$ GeV as the maximum value because it is constraint by the mass of the $m_{Z'} = (0.5-3)$ TeV, which is proportional to v_χ [3]

In our numerical analysis we have not considered the

contribution of gluon-gluon fusion because it is, at least, four orders of magnitude smaller than that of Drell-Yan process. To turn this work more complete we are introducing the width of $H^{\pm\pm} \rightarrow Uh^0$

$$\Gamma_{H^{\pm\pm} \rightarrow U^{\pm\pm} h^0} = \frac{AK}{32\pi m_{H^{++}}} \left(\frac{m_{H^{\pm\pm}}^4}{2m_{U^{\pm\pm}}^2} - \frac{m_{H^{\pm\pm}}^2 m_{h^0}^2}{m_{U^{\pm\pm}}^2} - m_{H^{\pm\pm}}^2 + \frac{m_{U^{\pm\pm}}^2}{2} + \frac{m_{h^0}^4}{2m_{U^{\pm\pm}}^2} - \frac{m_{h^0}^2}{2} \right), \quad (7)$$

where

$$A = \sqrt{\left[1 - \left(\frac{m_{U^{\pm\pm}} + m_{h^0}}{m_{H^{\pm\pm}}}\right)^2\right] \left[1 - \left(\frac{m_{U^{\pm\pm}} - m_{h^0}}{m_{H^{\pm\pm}}}\right)^2\right]}, \quad (8a)$$

$$K = \frac{ie u}{2s_W \sqrt{2(u^2 + w^2)}}. \quad (8b)$$




FIG. 6: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -1.2$ for the bosonic sector.

Considering that the expected integrated luminosity for the LHC will be of order of 3×10^5 pb $^{-1}$ /yr then the statistics give a total of $\simeq 3.9 \times 10^6$ events per year for Drell-Yan process, if we take the mass of the Higgs boson $m_{H^{\pm\pm}} = 500$ GeV, $v_\chi = 1500$ GeV, $m_{H_2^\pm}^\pm = 1163.7$ GeV, $m_h = 2229.9$ GeV and $\lambda_9 = -0.8$. Considering that the signal for $H^{\pm\pm}$ production will be e^-P^- and




FIG. 7: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -1.2$ for the $H^{\pm\pm} \rightarrow V^\pm H_1^\pm$ and $H^{\pm\pm} \rightarrow H_1^\pm H_2^\pm$.

e^+P^+ and taking into account that the branching ratios for both particles would be $B(H^{--} \rightarrow e^-P^-) = 3.6\%$ and $B(H^{++} \rightarrow e^+P^+) = 96.4\%$, see Fig. 3, we would have approximately 1.3×10^5 events per year for Drell-Yan process. Taking now the mass of the Higgs boson $m_{H^{\pm\pm}} = 700$ GeV, $v_\chi = 1500$ GeV, $m_{H_2^\pm}^\pm = 1223.6$ GeV,




FIG. 8: Total cross section for the process $pp \rightarrow H^{--} H^{++}$ as a function of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$




FIG. 9: Total cross section for the process $pp \rightarrow H^{--} H^{++}$ as a function of $m_{H^{\pm\pm}}$ for $\lambda_9 = -1.2$

$m_h = 2052.2$ GeV and the same value of λ_9 as above.

We then have a total of $\simeq 2.2 \times 10^6$ events per year for Drell-Yan process. Considering now the same signal as above, whose branching ratios are equal to $B(H^{--} \rightarrow e^- P^-) = 0.8\%$ and $B(H^{++} \rightarrow e^+ P^+) = 22\%$, we will have a total of approximately 3.9×10^3 events for Drell-Yan process. In Fig. 9 we show the cross section for the process $pp \rightarrow H^{++} H^{--}$ for the same parameter values of λ'_s , $v_\eta = 195$ GeV and other particle masses considered above except for the $\lambda_9 = -1.2$, which gives for $m_{H^{\pm\pm}} = 500$ GeV, $v_\chi = 1500$ GeV, the values of $m_{H_2^\pm}^\pm = 901.6$ GeV and $m_h = 2802.7$ GeV, in this case the statistics will be equal as in Fig. 8, because the cross section are nearly equal and also the branching ratios, where participate the same particles for the mass $m_{H^{\pm\pm}} = 500(700)$ GeV, are also equals. So from this analysis we conclude that varying the value of λ_9 from $-3 < \lambda_9 < 0$, where in our case we choose $\lambda_9 = -0.8$ and $\lambda_9 = -1.2$, and choosing the same signals, we obtain results nearly equal. Then we can conclude that we will have a very striking signal, the double-charged Higgs boson will deposit four times the ionization energy than the characteristic single-charged particle, that is, if we see this signal we will not only be seeing the double charged Higgs but also the heavy leptons. The main background for this signal, $pp \rightarrow H^{--} H^{++} \rightarrow e^- P^- (e^+ P^+)$, could come from the process $pp \rightarrow Z Z$ and another small background from the $pp \rightarrow W^- W^+ Z$, all these backgrounds can be eliminated (see Ref. [9]).

In summary, through this work, we have shown that in the context of the 3-3-1 model the signatures for double charged Higgs bosons can be significant in LHC collider. Our study indicates the possibility of obtaining a clear signal of these new particles with a satisfactory number of events.

TABLE IV: Numbers of events per year for doubly charged higgs Bosons with respect their masses and branching ratios. The total number of events per year for $m_{H^{++}} = 500$ GeV is 1.5×10^6 and for $m_{H^{++}} = 700$ GeV is 1.5×10^3 .

$BR(H^{--} \rightarrow e^- E^-)$	$BR(H^{++} \rightarrow e^+ E^+)$	$m_{H^{++}}^2$ (GeV)	Events/yr
3.6%	96.4%	500	5.5×10^4
0.8%	22%	700	1.6×10^3

I. ACKNOWLEDGMENTS

One of us (JECM) would like to thank Prof. R. O. Ramos for a careful reading of the manuscript and an-

other (N. V. C.) would like to thank the Instituto de Física Teórica, of UNESP, for the use of its facilities.

[1] V. Barger *et al.* Phys. Rev. D **26**, 28 (1982); R. N. Mohapatra and J. D. Vergados, Phys. Rev. Lett. **47**, 1713

(1981); G. B. Gelmini and M. Roncadelli, Phys. Lett. B,

- 99**, 411 (1981).
- [2] R.N. Mohapatra and P.B. Pal, *Massive Neutrinos in Physics and Astro-physics*, Singapore, Singapore: World Scientific, World Scientific lecture notes in physics, 41 (1991) 318 pp; R. N. Mohapatra and J.C. Pati, Phys. Rev. D **11** (1975) 566; G. Senjanovic and R. N. Mohapatra, Phys. Rev. D **12** (1975) 1502; T. Rizzo, Phys. Rev. D **25** (1982) 1355.
 - [3] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992); R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, *ibid* **47**, 4158 (1993).
 - [4] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48**, 2353 (1993).
 - [5] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48**, 5274 (1993).
 - [6] M. D. Tonasse, Phys. Lett. B **381**, 191 (1996).
 - [7] J. E. Cieza Montalvo, Nelson V. Cortez, J. Sa Borges and M. D. Tonasse, Nucl. Phys. B **756**, 1 (2006).
 - [8] W. -M. Yau *et al.* (Particle Data Group), J. Phys. G: Nucl. Part. Phys. **33**, 1 (2006).
 - [9] J. E. Cieza Montalvo and P. P. Queiróz Filho, Phys. Rev. D **66**, 055003 (2002).